



OPEN ACCESS JOURNAL



Check for updates

**Citation:** Mukesh Kumar, Sushil Prasad, Ravindra Kumar, Nandani Kumari, Soumen Naskar, Pankaj Kumar, Pawan Kumar Verma (2021). Effect of Generations and Different Agro Climatic Zones an Important Reproductive Trait in Black Bengal Goats. *Chemical Engineering*. v02i02, 13-20. <http://dx.doi.org/10.53709/CHE.2021.v02i02.003>

**DOI:**  
<http://dx.doi.org/10.53709/CHE.2021.v02i02.003>

**Corresponding Author:**  
Mukesh Kumar  
[mukesh2509.mk@gmail.com](mailto:mukesh2509.mk@gmail.com)

**Received on:** December 10, 2020  
**Revised on:** February 23, 2021  
**Accepted on:** June 03, 2021

**Copyright:** © 2021 Mukesh Kumar. Published under a [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) (CC BY 4.0) license.

## RESEARCH ARTICLE

# Effect of Generations and Different Agro Climatic Zones an Important Reproductive Trait in Black Bengal Goats

Mukesh Kumar<sup>1\*</sup>, Sushil Prasad<sup>1</sup>, Ravindra Kumar<sup>1</sup>, Nandani Kumari<sup>2</sup>, Soumen Naskar<sup>3</sup>, Pankaj Kumar<sup>4</sup>, Pawan Kumar Verma<sup>1</sup>

<sup>1</sup>Department of Livestock Production and Management, Ranchi Veterinary College, Bisra Agricultural University, Ranchi, Jharkhand 834006, India

<sup>2</sup>Department of Animal Genetics and Breeding, Ranchi Veterinary Collage, Bisra Agricultural University, Ranchi 834006, India

<sup>3</sup>Indian Council of Agricultural Research-Indian Institute of Agricultural Biotechnology, Namkum Ranchi 834010, India

<sup>4</sup>Livelihood Specialist, Wazirganj Gaya Bihar, India 834006, India

## ABSTRACT

Goat farming is a traditional occupation and an integral part of rural India's symbiotic system of agriculture and livestock production. For Jharkhand, it is a legacy breed due to superior quality meat and skin. Black Bengal goat farming plays an important role in improving the present investigations were carried out on Black Bengal goats maintained at AICRP BAU. Three agro-climatic zones, namely central and Northeastern plateau (IV), Ranchi and Deoghar, were selected. Similarly, from the Western plateau (V), East-Singbhum, and from South Eastern plateau (VI), Lohardhaga were respectively selected for the study. A cluster of four villages each from the four Districts above were taken, namely Chamguru, Palajori, Barabanki, and Tiko. Twenty farmers were each taken from each of the clusters, making a total of eighty farmers. The study was conducted on 80 farmers, 1600 Doe, 2459 kidding and their 4256 kids (male 2223 kids and female 2027 kids) during a period of six years maintained under All India Coordinated Research Project (AICRP) from 2014 to 2020 on Goat Improvement were observed. The average age at first estrus (days) for different agro-climatic zones and villages were observed to be  $237.28 \pm 1.28$ ,  $238.9 \pm 1.25$ ,  $240.2 \pm 1.2$ , and  $240.18 \pm 1.1$  days for the villages Chamguru, Palajori, Tiko, and Barabanki, respectively. Pooled overall age at first estrus was observed to be  $239.14 \pm 0.06$  for all the districts taken together. Age at first estrus (days) was observed to be highest for the village Tiko ( $240.2 \pm 1.2$ ) followed by Barabanki ( $240.18 \pm 1.1$ ), Palajori ( $238.9 \pm 1.25$ ), Chamguru ( $237.28 \pm 1.28$ ), and the effect of the village was significant ( $P < 0.05$ ).

**Keywords:** Colloidal processing, Perovskite Quantum dots, Solar energy materials, Metal halide, Electron microscope

## INTRODUCTION

Goat rearing is an essential part of the lives of poor farmers in Jharkhand and of the rural economy. Goats are often referred to as the poor man's cow and contribute to the farmers' livelihoods, especially for the women in the household. Despite its rising prices, goats are also an essential part of society and culture, especially for the tribal farmers. It being a slice of meat without any social or religious taboo is the preferred meat during marriages and festivals despite its ever-rising prices. At the global level, India stands second in goat population and accounts for 26% of global meat production. The small ruminant population of India, about 222.6 million heads of which about 148.88 million in goat (census 2019) in Jharkhand goat population 9.12 million [13] GDP is minimal (0.61) as compared with other sectors. The Goat population in the country in 2019 is 148.88 million showing an increase of 10.1%. Total milk production in India is 187.75 million tones and 4% of the total milk production [13] and goat meat production 13.53% in Jharkhand goat meat production is 21.68 million tones. Meat production in India increased by 2.88% annually from (livestock census 2019), but demand continues to exceed supply in the

country as well as in Jharkhand. The development of the goat sector, therefore, holds excellent opportunity for farmers and to contribute in a more significant manner to their household income. Reproductive performance is one of the main determinants of productivity as well as profitability. Proper selection, breeding, and management to improvise important reproductive and productive traits are essential for profit in Black Bengal goat production. Interaction of genes and environment plays a crucial role besides the genotype and environment effect in deciding the phenotype of any animal. Parity wise Reproductive efficiency as such can be measured and expressed as the kidding rate, kidding interval, litter size at birth, and litter weight at birth. Black Bengal goat is a popular, resilient, and prolific breed of goat highly preferred among the goat rearers. Thus the objective of the present study was:-To evaluate the reproductive and production performance and production potential of Black Bengal goats.

## MATERIAL AND METHODS

The study was conducted on 80 farmers, 1600 does, 2499 kidding, and their 4256 kids (male 2223 kids and female 2027 kids) during period six (6) years maintained under All India Coordinated Research Project on Goat Improvement [AICRP], Black Bengal field unit, BAU. The present investigations were carried out on Black Bengal goat maintained at AICRP BAU. Three agro-climatic zones, namely central and Northeastern plateau (IV), Ranchi and Deoghar were selected. Similarly, from the Western plateau (V), East-Singbhum, and from South Eastern plateau (VI), Lohardhaga was respectively selected for the study. A cluster of four villages each from the four Districts above were taken, namely Chamguru, Palajori, Barabanki, and Tiko. Twenty farmers were each taken from each of the clusters making a total of eighty farmers. The study was conducted on 80 farmers, 1600 Doe, 2499 kidding and their 4256 kids (male 2223 kids and female 2027 kids) during a period of six years maintained under All India Coordinated Research Project (AICRP) on Goat Improvement, Black Bengal field unit, BAU from 2014 to 2020. Farmers having a minimum number of

twenty goats were selected randomly. Reproductive and productive traits were studied in the experimental animals. For molecular characterization, ten goats were selected from each cluster; the data was recorded and collected from animals of three agro-climatic zones of four BAU cluster villages of Jharkhand. Cropped Area 851.05 thousand hectares Irrigated Area to Cropped Area 11.40%, Altitude, 275-670 m, average rainfall: 1270mm,, temperature 4.8-44.2°C. Important Crops: Rice, Wheat, Maize, Ragi, Potato, Mustard-Rapeseeds Gram and Pulses. Statistical analysis of the data collected was done using WINSTAT trial version. One way ANOVA was the Parametric test of hypothesis testing applied to see if the effect of the district was significant or not and to see the year-wise variation on different economic traits. The results have been tabulated in Table 2.

## RESULTS AND DISCUSSION

The reproductive performances of black Bengal goats viz age at first estrus, age at first service, age at first kidding, kidding interval, and litter size at birth, litter weight at birth were obtained. Various reproductive traits, namely, age at first estrus, age at first service, age at first kidding, kidding interval, litter size, the number of kid produced for a total of eighty farmers, and 1600 does have been reported for four villages under study.

### Effect of Agro-climatic zone on reproductive performance of black Bengal goat.

Average reproduction performance of Black Bengal goats belonging to the three different agro-climatic zones and four villages, namely Central and North Eastern Plateau (Chamguru and Palajori), Western Plateau (Tiko), and Southern Plateau (Barabanki) were observed. The average Age at first estrus(days) for different agro-climatic zones and villages were observed to be  $237.28 \pm 1.28$ ,  $238.9 \pm 1.25$ ,  $240.2 \pm 1.2$ , and  $240.18 \pm 1.1$  days for the villages Chamguru, Palajori, Tiko, and Barabanki, respectively Pooled overall age at first estrus were observed to be  $239.14 \pm 0.06$  for all the districts taken together. Age at first estrus (days) was

**Table 2.** Agro climatic zone and cluster village wise selected farmers and animals Population

Agro –climatic zone	Name of cluster Districts and village	Total Number of farmers	Total Number of goats	Total Number of kids		Number of total kids
				Male	Female	
Central and north eastern plateau	Ranchi Chamguru	20	400	569	501	1070
	Deoghar Palajori	20	400	475	555	1030
Western plateau	Lohardaga Tiko	20	400	499	489	986
Southern plateau	East-singhbhum Barabanki	20	400	683	487	1170
	Total	80	1600	2223	2027	4256

observed to be highest for the village Tiko ( $240.2 \pm 1.2$ ) followed by Barabanki ( $240.18 \pm 1.1$ ), Palajori ( $238.9 \pm 1.25$ ), and Chamguru ( $237.28 \pm 1.28$ ), and the effect of the village was significant ( $P < 0.05$ ). Table 3.1 Ali *et al.*, (2016) reported by Average reproductive performance of Black Bengal goats under farming conditions and management.

Average Age at first service (days) during different (three) agro-climatic zones and the four villages were observed to be  $257.23 \pm 1.2$ ,  $257.95 \pm 1.2$ ,  $240.2 \pm 1.2$  and  $257.71 \pm 1.1$  days for Chamguru, Palajori, Tiko and Barabanki, respectively. Overall Age at first service (days) was observed to be  $257.74 \pm 0.61$  for all the villages taken together. A significant difference ( $P < 0.05$ ) for age at first service (days) was observed among villages Tiko ( $240.2 \pm 1.2$ ) followed by Palajori ( $257.95 \pm 1.2$ ), Barabanki ( $257.71 \pm 1.1$ ), and Chamguru ( $257.23 \pm 1.2$ ). Table 3.1 The age at first service in Bengal goat vary from 200 to 240 days. Das *et al.*, 2008 [12] and Dhara *et al.*, 2008 [14] and a higher range (296 to 332 days) This study recorded the age of effective service for Bengal goat as 261.35 days with a range of 113 to 483 days.

Average Age at first Kidding (days) for different (three) agro-climatic zones and (four) villages were observed to be  $423.49 \pm 1.2$ ,  $421.31 \pm 1.2$ ,  $420.16 \pm 1.2$ , and  $415.41 \pm 1.1$  days for the village Chamguru, Palajori, Tiko, and Barabanki, respectively. Overall Age at first kidding (days) was observed to be  $420.00 \pm 0.00$  for all the villages taken together. A significant difference ( $P < 0.05$ ) for age at first Kidding (days) were observed among village Chamguru ( $423.49 \pm 1.2$ ) followed by Palajori ( $421.31 \pm 1.2$ ), Tiko ( $420.16 \pm 1.2$ ) and Barabanki ( $415.41 \pm 1.1$ ) days respectively. Table 3.1 Faruque *et al.* (2010) [16] calculated the average age at first kidding to be  $283.83 \pm 31.16$  day's and  $370.26 \pm 25.48$  days for intensive and semi intensive system. Halim *et al.* (2011) [21] reported the age at first kidding for Black Bengal goat to be  $361.83 \pm 11.29$  days. Bhowmik *et al.* [8] has reported the age at first kidding for Jamunapari, Black Bengal, and crossbred goats to be  $534 \pm 24.58$ ,  $368.12 \pm 16.96$  and  $471.25 \pm 21.25$  days, respectively. Hasan *et al.* (2015) [20] reported the age at first kidding to be  $448.26 \pm 25.48$  days under semi-intensive conditions, whereas under extensive conditions, it was recorded to be  $450.07 \pm 22.43$  days.

Average Kidding interval (days) during different (three) agro-climatic and (four) villages were observed to be  $241.4 \pm 1.2$ ,  $240.68 \pm 1.2$ ,  $240.28 \pm 1.2$ , and  $241.21 \pm 1.1$  days during village Chamguru, Palajori, Tiko, and Barabanki, respectively. Overall, Kidding interval days were observed to be  $240.9 \pm 0.61$  respectively. Significant different highest ( $P < 0.05$ ) Kidding interval days were observed to be among village Chamguru ( $241.4 \pm 1.2$ ) followed by Barabanki ( $241.21 \pm 1.1$ ),

Palajori ( $240.68 \pm 1.2$ ), and Tiko ( $240.28 \pm 1.2$ ) days respectively. However, significant differences ( $P < 0.05$ ) were observed among village Table 3.2. The kidding interval of animal is breed-specific but it varies due to management condition in which they prevail. Most of the previous workers reported the kidding interval for Black Bengal goat Faruque *et al.* [16] found that the average kidding interval was  $181.23 \pm 4.55$  days for Black Bengal goats. Mamabolo *et al.* [28] reported the average kidding interval of goats in South Africa at village level to be 145-148 days. Bhowmik *et al.* [8] has reported the first kidding interval for Jamunapari, Black Bengal, and crossbred goats to be  $224 \pm 14.42$ ,  $181.76 \pm 15.81$  and  $199.17 \pm 21.71$  days respectively [20] found that the kidding interval was  $178.23 \pm 50$  days under semi-intensive conditions and  $190.20 \pm 20$  under an extensive system.

Average Litter size at birth (nos.) for the different (three) agro-climatic and (four) villages were observed to be  $2.24 \pm 0.00$ ,  $2.30 \pm 0.00$ ,  $2.32 \pm 0.00$  and  $2.16 \pm 0.00$  days for the villages Chamguru, Palajori, Tiko, and Barabanki, respectively. Overall, Litter size at birth (nos.) was found out to be  $2.25 \pm 0.00$  for all the villages taken together. A significant difference ( $P < 0.05$ ) was found on litter size at birth (nos.) among the villages Tiko ( $2.32 \pm 0.00$ ), Palajori ( $2.30 \pm 0.00$ ), Chamguru ( $2.24 \pm 0.00$ ), and Barabanki ( $2.16 \pm 0.00$ ). Table 3.1 Hasan *et al.* (2015) [20] found that the average birth weight of kids was  $1.28 \pm 0.11$  kg and  $1.25 \pm 0.10$  kg under semi-intensive and an extensive system of management, respectively and Paul *et al.* (2014) reported the average birth weight of Black Bengal kids in NBF, Modhupur and Dimla were  $1.09 \pm 0.027$  kg,  $1.09 \pm 0.025$  and  $1.10 \pm 0.027$  kg respectively, Bhowmik *et al.* (2014) [8] has reported that the average birth weight of Jamunapari, Black Bengal and cross bred goats were  $1.53 \pm 0.22$  kg,  $0.89 \pm 0.09$  and  $1.26 \pm 0.20$  kg respectively.

Average Litter weight at birth (Kg) for different (three) agro-climatic zones and the (four) villages were observed to be  $3.28 \pm 1.2$ ,  $3.27 \pm 1.2$ ,  $3.26 \pm 1.2$  and  $3.27 \pm 1.2$  days for the villages Chamguru, Palajori, Tiko, and Barabanki, respectively. Overall, Litter weight at birth (Kg) was found out to be  $3.27 \pm 0.61$  for all the cluster villages pooled together. Significant ( $P < 0.05$ ) effect on litter weight at birth (Kg) was found between the four villages, namely Chamguru ( $3.28 \pm 1.2$ ), Palajori ( $3.27 \pm 1.21$ ), Barabanki ( $3.27 \pm 1.2$ ), and Tiko ( $3.26 \pm 1.2$ ). (Table 3.1 and fig 3.1) They also recorded that the gains of triplets kids were the lowest but steady for 0-3 and 3-6 months, whereas gains of single and twin kids sharply declined from 6-9 months. The significant effect of the village on 6-month body weight indicates that the genetic pool and management practices of village 2 are poor. However, the 6-month body wt. of Black Bengal kids recorded in this study as

**Table 3.1.** Effect of Agro-climatic zone on reproductive performance of Black Bengal goat

Parameter/village	Central and North Eastern Plateau		Western Plateau	Southern Plateau	Overall
	Chamguru	Palajori	Tiko SE	Barabanki	
Age at first estrus(day)	237.28±1.28 <sup>a</sup> (400)	238.9±1.25 <sup>b</sup> (400)	240.2±1.2 <sup>c</sup> (400)	240.18±1.1 <sup>c</sup> (400)	239.14±0.61 (1600)
Age at first service (day)	257.23±1.2 <sup>a</sup> (400)	257.95±1.2 <sup>a</sup> (400)	258.08±1.2 <sup>b</sup> (400)	257.71±1.1 <sup>a</sup> (400)	257.74±0.61 (1600)
Age at first Kidding(-day)	423.49±1.2 <sup>c</sup> (400)	421.31±1.2 <sup>ab</sup> (400)	420.16±1.2 <sup>b</sup> (400)	415.41±1.1 <sup>a</sup> (400)	420.00±0.00 (1600)
Kidding interval(day)	241.4±1.2 <sup>b</sup> (400)	240.68±1.2 <sup>a</sup> (400)	240.28±1.2 <sup>a</sup> (400)	241.21±1.1 <sup>b</sup> (400)	240.9±0.61 (1600)
Litter size at birth (No)	2.24±0.00 <sup>b</sup> (559)	2.30±0.00 <sup>c</sup> (564)	2.32±0.00 <sup>c</sup> (685)	2.16±0.00 <sup>a</sup> (651)	2.25±0.00 (2459)
Litter weight at birth (Kg)	3.28±1.2 <sup>c</sup> (559)	3.27±1.21.2 <sup>b</sup> (564)	3.26±1.2 <sup>a</sup> (685)	3.27±1.2 <sup>b</sup> (651)	3.27±0.61 (2459)

Figure in parentheses indicate the number of dams

\* P < 0.05, \*\* P < 0.01, NS = Non-Significant

7.96 ± 0.38 kg was within the range of 6.82 kg to 10.59 kg as observed by earlier workers at different times and different places Hasan et al. (2015) [20] found that average litter weight of kids was 4.28±0.11kg and 5.25±0.10kg under semi-intensive and an extensive system of management respectively. Paul et al. (2014) [35] reported the average birth weight of Black Bengal kids in NBF, Modhupur, and Dimla were 3.09 ± 0.027 kg, 5.09± 0.025 and 5.10± 0.027kg, respectively. Bhowmik et al. (2014) [8] has reported that the average litter weight at birth of Jamunapari, Black Bengal, and crossbred goats were 1.53±0.22kg, 0.89±0.09, and 1.26±0.20 kg, respectively.

### Effect of Black Bengal (goats) Dam performance up to 6<sup>th</sup> Parity

The data for important genetic traits like age of kidding, kidding interval, litter size, and litter weight was collected and tabulated, and it was observed that the Black Bengal goats had the genetic capability to produce first kid as early as 1.13 years and by 4.39 years it completed 6<sup>th</sup> kidding. The kidding interval ranged from 171.85 to 222.30 in parity one to 216 days in the 6<sup>th</sup> parity, and the variation in the kidding interval was significant (P<0.05) w.r.t the number of parity. The average litter size in Black Bengal goats in different parity ranged from 2.21 to 2.31, and litter weight ranged from 3.26 to 3.39. The effect of parity was significant (p<0.05) on both the traits, namely litter size at birth and litter weight at birth. (Table-3.2 reported that the productive life period is almost half compared to the average life period, although they studied in cattle. As per Alam (2003) [5] the average fertile life period of Black Bengal doe in Bangladesh was 7–8 years and 7–10 years, respectively. Faruque

et al. (2010) [16], Hasan et al. (2015) [20], and Paul et al. (2014) [35] reported a significant effect of parity on the birth weight of kids. Average litter size (1.76±0.08) of Black Bengal goats is almost the same as the results of Amin et al. (2014) [7], who reported 1.9±0.56, 1.86, 1.61±0.02 litter size, respectively. Variation in litter size may be due to the interaction of genetics and environment, status and age of the dam, parity. Abraham et al. (2019)[1] reported that the effect of genetic and non-genetic factors on the reproductive performance of Sirohi and its crosses with the Beetal (half-bred) in the semi-arid conditions of Rajasthan were studied. The crossbred has significantly less kidding interval than Sirohi does. There was no effect of season on the reproductive traits studied. Misra and Sinha (2001) [29] reported the average kidding interval in Black Bengal as 8.80±0.03 months. Hassan et al. (2007) [18] reported the average interval in Black Bengal and crossbred goats as 179±20 and 270±22 days, respectively, and the differences are statistically significant (P<0.01). Hasan et al. (2015) [20] found the non-significant effect of parity on kidding interval. Akhtar et al. (2006) [4] reported that the mean body weight of Black Bengal at 3 months age were 4.99±0.15, 4.64±0.33, 4.07±0.42 and 4.38±0.13, 4.39±0.27, 3.99±0.47 kg respectively for three generations of selected and control groups. Jimenez et al. (2009) in Serrana Transmontano kids, in Nigeria Sahelian goats, have reported significant effect of parity of dam on body weight of kids


### Effect of different Generation wise reproductive parameters of Black Bengal Goat.

A study was conducted to see the effect of generation on different reproductive parameters. Data collected


**Table 3.2.** Effect of Black Bengal (goats) Dam performance up to 6<sup>th</sup> Parity

Parity/parameter	Age at kidding (Years)	Kidding interval (day)	Litter size (No)	Litter weight(Kg)
1 <sup>st</sup> parity	1.13±0.02 (400)		2.21±0.14 <sup>ab</sup> (181)	3.29±0.17 <sup>d</sup> (181)
2 <sup>nd</sup> parity	1.85±0.06 (309)	216.30±0.00 <sup>d</sup> (309)	2.31±0.14 <sup>b</sup> (134)	3.26±0.18 <sup>ab</sup> (134)
3 <sup>rd</sup> parity	2.50±0.00 (263)	195.25±0.60 <sup>c</sup> (263)	2.16±0.00 <sup>a</sup> (122)	3.27±0.19 <sup>c</sup> (122)
4 <sup>th</sup> parity	3.11±0.04 (185)	183.91±0.6 <sup>b</sup> (185)	2.26±0.16 <sup>ab</sup> (80)	3.24±0.00 <sup>a</sup> (80)
5 <sup>th</sup> parity	3.85±0.03 (107)	222.30±0.60 <sup>e</sup> (107)	2.31±0.16 <sup>ab</sup> (47)	3.24±0.21 <sup>a</sup> (47)
6 <sup>th</sup> parity	4.39±0.06 (69)	171.83±0.74 <sup>a</sup> (69)	2.17±0.18 <sup>ab</sup> (32)	3.27±0.23 <sup>c</sup> (32)
Overall			2.23±0.12 (591)	3.26±0.16 (591)

Fig:- in parentheses indicate number of dams/litter  
 \* P < 0.05, \*\* P < 0.01, NS = Non-Significant




**Fig 3.1.** Effect of Agro-climatic zone wise reproductive performance of Black Bengal goat.



**Fig 3.2.** Effect of Black Bengal (goats) Dam performance up to 6<sup>th</sup> Parity

for various economic traits like Age at first estrus (day), Age at first service (day), Age at first Kidding (day), Kidding interval (day), Litter size (No), and Litter weight (Kg) for six generations in the experimental Black Bengal goats. Age at first estrus (day), was to found to the significantly lowest in first-generation followed by 4<sup>th</sup> generation. The maximum value was found in the fifth generation. The average age at first



**Fig:- 3.3.** Effect of different Generation wise reproductive parameter of black Bengal Goat.

estrus (day) was 239.14±0.61 days. Similarly, age at first service (day) was found to the lowest in first generation and the highest Age at first service (days) was reported in 2<sup>nd</sup> generation with a mean value 257.47±00 days and 258.12±0.55 days, respectively. The average mean value of age at first service (days) was 257.74±0.64, respectively. Age at first kidding (days) was observed to the 424.67±2.5, 420.03±2.53, 415.33±00, 421.83±2.5, 415.34±2.5, and 421.78±2.5 days respectively for the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> generation respectively. The age at first service in Bengal goat varies from 200 to 240 days [42; Das et al., (2008) [12] and Dhara et al., (2008) [14], and a higher range (296 to 332 days) has also been reported by this study recorded the age of effective service for Bengal goat as 261.35 days with a range of 113 to 483 days.

The average value was found out to the 420.00±0.61 days. The kidding interval was reported to be 424.67±2.5, 420.03±2.53, 415.33±00, 421.83±2.5, 415.34±2.5 and 421.78±2.5 for the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> generation respectively. The kidding interval of

**Table 3.3.** Effect of different Generation wise reproductive parameters of Black Bengal Goat

Parameter/ generation	1 <sup>st</sup> G Mean± SE	2 <sup>nd</sup> G Mean± SE	3 <sup>rd</sup> G Mean± SE	4 <sup>th</sup> G Mean± SE	5 <sup>th</sup> G Mean± SE	6 <sup>th</sup> G Mean± SE	Overall Mean± SE
Age at first es- trus(day)	237.22±0.00 <sup>a</sup> (58)	240.17±1.4 <sup>b</sup> (38)	240.17±1.4 <sup>b</sup> (36)	238.85±1.4 <sup>b</sup> (32)	240.24±1.4 <sup>b</sup> (25)	238.86±1.4 <sup>b</sup> (12)	239.14±0.61 (201)
Age at first ser- vice (day)	257.47±0.00 <sup>a</sup> (58)	258.12±0.55 <sup>b</sup> (38)	257.66±0.55 <sup>ab</sup> (36)	258.03±0.55 <sup>ab</sup> (32)	258.09±0.55 <sup>b</sup> (25)	258.03±3.5 <sup>b</sup> (12)	257.74±0.61 (201)
Age at first Kid- ding(day)	424.67±2.5 <sup>c</sup> (58)	420.03±2.53 <sup>b</sup> (38)	415.33±0.00 <sup>a</sup> (36)	421.83±2.5 <sup>b</sup> (32)	415.34±2.5 <sup>a</sup> (25)	421.78±2.5 <sup>b</sup> (12)	420.00±0.61 (201)
Kidding inter- val(day)	241.79±0.5 <sup>c</sup> (58)	240.26±0.00 <sup>a</sup> (38)	241.26±0.51 <sup>b</sup> (36)	241.01±0.51 <sup>b</sup> (32)	241.25±0.51 <sup>b</sup> (25)	241.01±0.51 <sup>b</sup> (12)	240.9±0.61 (201)
Litter size (No)	2.21±0.12 <sup>ab</sup> (122)	2.32±0.12 <sup>b</sup> (80)	2.16±0.00 <sup>a</sup> (68)	2.25±0.12 <sup>ab</sup> (61)	2.31±0.12 <sup>b</sup> (45)	2.16±0.12 <sup>a</sup> (18)	2.25±0.61 (394)
Litter weight (Kg)	3.29±0.14 <sup>a</sup> (122)	3.27±0.14 <sup>a</sup> (80)	3.27±1.4 <sup>a</sup> (68)	3.25±0.00 <sup>a</sup> (61)	3.26±0.14 <sup>a</sup> (45)	3.27±0.14 <sup>a</sup> (18)	3.27±0.61 (394)

Fig:- in parentheses indicate the number of dams/litter

\* P < 0.05, \*\* P < 0.01, NS = Non-Significant

animals is breeding specific but it is varied chiefly due to management condition in which they prevail. Most of the previous workers reported the kidding interval for Black Bengal goat 176 – 240 days Misra and Sinha, (2001) [29] Das et al., (2008) [12] and Dhara et al., (2008) [14]. This result confirms the kidding interval of Black Bengal goat as 230–240 days even in farmer's conditions. Abraham et al. (2019) [1] found the kidding interval of does under semi-intensive system (233.8±13.12 days) than those of the does manage under an extensive system (242.8±32.66 days).

The average litter size at birth (kg) of the six-generation taken together was 2.25 ±0.06 in 6<sup>th</sup> generation and 2.21±0.12, 2.32±0.12, 2.16±0.00, 2.25±0.12, 2.31±0.12, and 2.16±0.12 in the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> generation respectively. They also recorded that the gains of triplets kids were the lowest but steady for 0 – 3 and 3 – 6 months whereas gains of single and twin kids sharply declined from 6 – 9 months. The significant effect of the village on 6-month body weight indicates that the genetic pool and/or management of village 2 are poor. However, the 6-month body wt. of Black Bengal kids recorded in this study as 7.96 ± 0.38 kg was within the range of 6.82 kg to 10.59 kg as observed by earlier workers at different times and different place average litter size (1.76±0.08) of Black Bengal goats which is almost same with the results of Amin et al. (2014) [7] and Paul et al. (2014) [35] who reported 1.9±0.56, 1.86, 1.61±0.02 litter size, respectively. Variation in litter size may be due to the interaction of genetics and environment, status and age of the dam, parity.

The average litter weight at birth (kg) of the six generations taken together was 3.27±0.61kg for the and the litter weight (kg) was observed to be 3.29±0.14, 3.27±0.14, 3.27±1.4, 3.25±0.00, 3.26±0.14 and 3.27±0.14 in the 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> generation respectively. They also recorded that the gains of triplets kids were the lowest but steady for 0 – 3 and 3 – 6 months whereas gains of single and twin kids sharply declined from 6 – 9 months. The significant effect of the village on 6-month body wt. indicates that the genetic pool and/or management of village 2 are poor. However, the 6-month body wt. of Black Bengal kids recorded in this study as 7.96 ± 0.38 kg was within the range of 6.82 kg to 10.59 kg as observed by earlier workers at different times and different place average litter weight of Black Bengal goats which is almost same with the results of Amin et al. (2000) [7] who reported 1.9±0.56, 1.86, 1.61±0.02 litter size, and litter weight respectively. Variation in litter size may be due to the interaction of genetics and environment, status and age of the dam, parity. Mishra and Sinha (2001) [29] reported that the effect of genetic and non-genetic factors on the reproductive performance of Sirohi and its crosses with the Beetal (half-bred) in the semi-arid conditions of Rajasthan were studied. The crossbred has significantly less kidding interval than Sirohi does. There was no effect of season on the reproductive traits studied. Abraham et al. (2019) [1] reported the average kidding interval in Black Bengal as 8.80±0.03 months. [18] reported the average interval in Black Bengal and crossbred goats as 179±20 and 270±22 days, respectively, and the differences are statistically significant (P<0.01). Hasan et al. (2015) found the non-significant effect of parity on kidding interval

## CONCLUSIONS

Black Bengal goats exhibit good performance records with respect to all reproductive and economic traits in the current research, so more farmers in the Jharkhand region of India should be encouraged to keep this breed of goat for their economic upliftment.

The effect of generation on litter weight was found to be non-significant. However, no definite increase or decrease in the value of any of the six economic traits was found with increasing parity from 1<sup>st</sup> to 6<sup>th</sup> generation.

## REFERENCE

- [1] Abraham H, S Gizaw and M Urge (2019). Growth and reproductive performances of Begait goat under semi intensive and extensive management in western Tigray, north Ethiopia. *Livestock Research for Rural Development*, 31. Article32. <http://www.lrrd.org/lrrd31/3/hagsa31032.html>.
- [2] Ali M, S Husain, S Bhuiyan, M Islam and M Haque (2016). Genetic evaluation of production potential of Black Bengal goat using frozen semen under community based rearing system. *Bangladesh Journal of Animal Science* 45(2):52-59.
- [3] Adodara RJ and N Jacob (2013). Climate change: Effect on growth of animals. *Basic Research Journal of Agricultural Science and Review* 2(4):80-90.
- [4] Akhtar, F., Islam, A.B.M.M. and Amin, M.R. 2006. Effect of selection for growth on production performance in Black Bengal goat. *Pakistan Journal of Biological Sciences* 9(2): 182-185.
- [5] Alam, F.M. (2003). Rc:Black Bengal goats breeds. *Agriscap*. July 31.01:40:00
- [6] Akhtar A (2018). Characterization and in situ improvement of Black Bengal goat at a community driven breeding program in Bangladesh. PhD Thesis. Department of Animal Breeding and Genetics, Bangladesh Agricultural University.
- [7] Amin MR (2014). Black Bengal – a promising goat genetic resource of Bangladesh, Joint FAO and IEA Program, International Atomic Energy Agency (2013-2014), P.O. Box 100, Wagramer Strasse 5, A-1400 Vienna, Austria.
- [8] Bhowmik N, MM Mia, MM Rahman and S Islam (2014). Preliminary study on productive and reproductive performances of Jamunapari, Black Bengal and crossbred goats at Chittagong region of Bangladesh. *Iranian Journal of applied Animal Science* 4:89-93.
- [9] Bushara I and Abu MMAA Nikhaila (2012). Productivity performance of Taggar female kids under grazing condition. *Journal of Animal Production Advances* 2(1):74-79.
- [10] Braham H, S Gizaw and M Urge (2019). Growth and reproductive performances of Begait goat under semi intensive and extensive management in western Tigray, north Ethiopia. *Livestock Research for Rural Development*, 31
- [11] Chowdhury SD (2015). Study on goat value chain in Bangladesh.
- [12] Das, A.K., Manna, M.K., Sahoo, A.K., Deb, R. and Paul, R.K. (2008) Studies on reproductive traits of Bengal goats in a part of West Bengal. *Indian journal of Small ruminants*. 14(2): 259-261.
- [13] Department of Animal Husbandry & Dairying releases 20th Livestock Census; Total Livestock population increases 4.6% over Census-2012, Increases to 535.78 million Posted On: 16 OCT 2019 6:26PM by PIB Delhi.
- [14] Dhara, K.C., Ray, N., Roy, S., Samanta, A.K. and Senapati, P.K. (2008). Improvement of Reproductive Performance of Black Bengal Goat through Selection under Field Condition. *J. Anim. Vet. Adv.*, 7 (5): 599-603.
- [15] DLS (2019). Livestock economy at a glance. [dls.portal.gov.bd>files>page>Updated Livestock Economy \(2018-2019\)](http://dls.portal.gov.bd/files/page/Updated%20Livestock%20Economy%20(2018-2019))
- [16] Faruque, S., Chowdhury, S.A., Siddiquee, N.U. and Afroz, M.A. (2010). performance of genetic parameters of economically important traits of Black Bengal goat. *J. Bangladesh Agril. Univ.* 8(1): 67-78.
- [17] Genetic Resource. (2008). DARE / ICAR Annual report 2007-2008. pp.17.
- [18] Hassan, M.M., Mahmud, S.M.N., Islam, S.K.M.A. and Maizi, O.F. (2007). A comparative study on Reproductive performance and productivity of the Black Bengal and crossbred goat at Atrai, Bangladesh. *Univ. J. Rajshahi Univ.* vol. 26, pp. 55-57.
- [19] Hasan MJ, Ahmed JU and Alam MM (2014). Reproductive performances of Black Bengal goat under semi-intensive and extensive conditions at rural areas in Bangladesh. *Journal of Advance Veterinary and Animal Research* 1: 196-200
- [20] Hasan, M.J., Ahmed, J.U., Alam, M.M., Mojumder, M.L.O. and Ali, M.S. (2015). Reproductive performance of Black Bengal goat under semi-intensive and extensive condition in Rajshahi district of Bangladesh. *Asian J. Med. Biol. Res.* 1(1),22-30.
- [21] Halim, M.A., Kashem, M.A., Mannan, A., Ahmed, SSU. and Hossain M.A. (2011). Reproductive and productive performance of Black Bengal goats reared by NGO beneficiaries under semi intensive system in Bangladesh. *International Journal of Natural science* 1(2).39-43.
- [22] Index Mundi(2019). Bangladesh- Rural population (% of total population).

- [23] Islam AFMF, S Nahar, TSJ Tanni, MA Hoque and AKFH Bhuiyan (2016). Breeding and production performance profile of Black Bengal goats in rural areas of Mymensingh in Bangladesh. *Bangladesh Journal of Animal Science* 45(2):46-51.
- [24] Islam MR, Amin MR, Kabir AKMA and Ahmed MU (2009). Comparative study between semi-intensive and scavenging production system on the performance of Black Bengal goat. *Journal of Bangladesh Agricultural University* 7: 79-86 05.
- [25] Jimenez- Badillo, M.R., Rodrigues, S., Sanudo, C., and Teixeira, C. (2009). Non-genetic factors affecting liveweight and daily weight in Serrana Transmontano Kids. Escola Superior Agraria de Braganca, centro de investigacao de Montanha. Apdo, 1172, 5300 Braganca, Portugal
- [26] Kumar A, Chae BJ, Bhuiyan AKFH, Sarker SC and Hossain MM (2018). Goat production system at Mymensingh district in Bangladesh. *Bangladesh Journal of Animal Science* 47: 13-20 06.
- [27] Moni IZ and Samad MA (2019). Evaluation of productive and reproductive performances of Black Bengal goats in Rajshahi Government Goat Development Farm in Bangladesh. *J. Vet. Med. OH Res.* 1(2): 201- 210 [DOI: 10.36111/jvmohr.2019.1(2).0012]
- [28] Mamabolo, M.J., Web, E.C., Preez, E.K. and Morns S.D. (2010). Reproductive status of goats in communal system in South Africa. Department of Animal and Wildlife Science, University of Pretoria, South Africa.
- [29] Mishra, S. K. and Sinha, R. (2001). Studies on the incidence of multiple birth and reproduction of Black Bengal goat under village conditions. *J. Interacad.* 5(2): 212-215.18
- [30] M Solaiman<sup>1</sup>, AS Apu<sup>1\*</sup>, MY Ali<sup>1</sup>, M Fakruzzaman<sup>2</sup> and MO Faruque<sup>1</sup> <sup>1</sup> Department of Animal Breeding and Genetics, Bangladesh Agricultural University. (2020) Impact of community based breeding program on breeding buck availability, growth and reproductive performance of Black Bengal goat Mymensingh-2202, Bangladesh; <sup>2</sup> Department of Genetics and Animal Breeding, Patuakhali Science and Technology University, Patuakhali, Bangladesh. *Bang. J. Anim. Sci.* 49 (1):13-21
- [31] MKI and J Khatun (2013). Different traits of Black Bengal goats under two feeding regime and fitting the Gompertz curve for prediction of weaning weight in the semi-scavenging system. *Indian Journal of Animal Research* 47: 498-503.
- [32] Monteiro A, Costa JM and Lima MJ (2017). Goat system productions: advantages and disadvantages to the animal, environment and farmer.
- [33] Miah G, Das A, Bilkis T, Momin MM, Uddin MA, Alim MA, Mahmud MS and Miazi OF (2016). Comparative study on productive and reproductive traits of Black Bengal and Jamunapari goats under semi-intensive condition. *Scientific Research Journal* 4: 1-7
- [34] Nandi D, Roy S, Bera S, Kesh SS and Samanta AK (2011). The rearing system of Black Bengal goat and their farmers in West Bengal, India. *Veterinary World* 4: 254-257.
- [35] Paul RC, Rahman ANMI, Debnath S and Khandoker MAMY (2014). Evaluation of productive and reproductive performance of Black Bengal goat. *Bangladesh Journal of Animal Science* 43: 104- 111 12.
- [36] Talukder MAI, Rahman MM, Alam MA and Hemayet MA (2016). Productive and reproductive performances of Brown Bengal goat (hilly goat) at research farm level. *Asian Journal of Medical and Biological Research* 2: 477-482 13.
- [37] Tribune (2018). UN: Bangladesh urban population will overtake rural population by 2010.
- [38] Roy S, SK Dash, SK Dhal, GD Nayak and AK Parida (2016). Genetic studies on reproductive performance of indigenous goats in Northern Odisha. *Exploratory Animal and Medical Research* 6(2):192-198.
- [39] R.K. (2008) Studies on reproductive traits of Bengal goats in a part of West Bengal. *Indian journal of Small ruminants*, 14(2): 259-261.
- [40] Siddiky NA (2017). Sustainable goat farming for livestock improvement in South Asia.
- [41] Uddin MN, Ahmed BS, Rashedunnabi AM, Uddin MMS, Hasan KMM and Baset MA (2014). Nutritional status of Black Bengal goat in government goat development farm, Sylhet. *Annals of Veterinary and Animal Science* 1: 97-100 15.
- [42] Zeshmarani, S., Dhara, K.C., Samanta, A.K., Samanta, R. and Majumder, S.C. (2007). Reproductive performance of Goats in Eastern and North-eastern India. *Livestock Research for Rural Development*, 19(8).